

FROBENIUS MANIFOLD STRUCTURE ON DOLBEAULT COHOMOLOGY AND MIRROR SYMMETRY

HUAI-DONG CAO & JIAN ZHOU

ABSTRACT. We construct a differential Gerstenhaber-Batalin-Vilkovisky algebra from Dolbeault complex of any closed Kähler manifold, and a Frobenius manifold structure on Dolbeault cohomology.

String theory leads to the mysterious Mirror Conjecture, see Yau [29] for the history. One of the mathematical predictions made by physicists based on this conjecture is the formula due to Candelas-de la Ossa-Green-Parkes [4] on the number of rational curves of any degree on a quintic in $\mathbb{CP}^4$. Recently, it has been proved by Lian-Liu-Yau [15].

The theory of quantum cohomology, also suggested by physicists, has lead to a better mathematical formulation of the Mirror Conjecture. As explained in Witten [28], there are two topological conformal field theories on a Calabi-Yau manifold X : A theory is independent of the complex structure of X , but depends on the Kähler form on X , while B theory is independent of the Kähler form of X , but depends on the complex structure of X . Vafa [25] explained how two quantum cohomology rings $\mathcal{R}$ and $\mathcal{R}'$ arise from these theories, and the notion of mirror symmetry could be translated into the equivalence of A theory on a Calabi-Yau manifold X with B theory on another Calabi-Yau manifold $\hat{X}$, called the mirror of X , such that the quantum ring $\mathcal{R}$ can be identified with $\mathcal{R}'$.

For a mathematical exposition in terms of variation of Hodge structures, see e.g. Morrison [19] or Bertin-Peters [2]. There are two natural Frobenius algebras on any Calabi-Yau n -fold,

$$A(X) = \bigoplus_{k=0}^n H^k(X, \Omega^k), \quad B(X) = \bigoplus_{k=0}^n H^k(X, \Omega^{-k}),$$

where Ω^{-k} is the sheaf of holomorphic sections to $\Lambda^k T^*X$. By Hodge theory, $B(X)$ can be identified with $H^n(X, \mathbb{C})$. By Bogomolov-Tian-Todorov theorem, there is a deformation of the complex structures on X parameterized by an open set in $H^1(X, \Omega^{-1})$. Therefore, one gets a family of Frobenius algebra structures on $H^n(X, \mathbb{C})$. Every Frobenius algebra structure can be characterized by a cubic polynomial Φ (in Physics literature, it is called the Yukawa coupling), so we get a family $\Phi_B(X)$ of cubic polynomials on $H^n(X, \mathbb{C})$ parameterized by an open set in $H^1(X, \Omega^{-1})$. An additional structure provided by algebraic geometry is the Gauss-Manin connection on this family. It is a flat connection with some extra properties. On the other hand, counting of rational curves provides a family $\Psi_A(X)$ of cubic polynomials on $A(X)$ parameterized by an open set in $H^1(X, \Omega^1)$. One version of Mirror Conjecture is the conjectural existence, for a Calabi-Yau 3-fold X , of another Calabi-Yau 3-fold $\hat{X}$, such that one can identify $B(X)$ with $A(\hat{X})$, $\Phi_B(X) = \Psi_A(\hat{X})$, and vice versa.

¹Both authors were supported in part by NSF

Witten [27] proposed that the associativity property of quantum cohomology can be encoded in WDVV (Witten-Dijkgraaf-Verlinde-Verlinde) equations (p.275, (3.14)). Dubrovin [7] studied family of Frobenius algebras which are parameterized by the underlying vector space, such that the family of structure cubic polynomials admits some symmetry, so that it becomes the third derivative of a single potential function. This potential function satisfies WDVV equations. Such a family will be called a *potential family*. He then introduced and studied the concept of Frobenius manifolds. See Dubrovin [8]. Ruan-Tian [21] gave a mathematical formulation of quantum cohomology (see also Liu [16] and McDuff-Salamon [17]). They also proved the potential property (WDVV equations) of the quantum cohomology, and gave the construction of a flat connection (suggested by Dubrovin's construction), which is the candidate for the mirror of Gauss-Manin connection. (In general, there is some convergence problem involved in their construction, except for the case of complete intersection Calabi-Yau manifolds. See Tian [23], §10.) Witten [28] suggested two kinds of extended moduli spaces, one containing the deformation space of the complex structure, the other containing the complexified Kähler cone. Kontsevich [11], besides reformulating Mirror Conjecture in a more general setting, proposed a construction of extended moduli space of complex structures. Recently, Barannikov-Kontsevich [1] constructed an extended formal moduli space for any Calabi-Yau manifold, as a generalization of Bogomolov-Tian-Todorov theorem. Furthermore, they showed that there is a structure of formal Frobenius manifold on it. Manin [18] has formulated a generalization to the construction of a formal Frobenius manifold from any differential Gerstenhaber-Batalin-Vilkovisky (dGBV) algebra with some mild conditions.

By Hodge theory (see e.g. Griffiths-Harris [10]), there is a decomposition

$$H^*(X, \mathbb{C}) \cong \bigoplus_{p,q} H^{p,q}(X),$$

where $H^{p,q}(X)$ is the Dolbeault cohomology. Furthermore, exterior product induces

$$H^{p,q}(X) \wedge H^{r,s}(X) \subset H^{p+r,q+s}(X).$$

In Cao-Zhou [5], the authors developed a quantum de Rham cohomology theory. In that theory, quantum Dolbeault cohomology can be defined such that it satisfies the usual properties of the ordinary Dolbeault cohomology. This suggests the construction of Frobenius manifold structure on Dolbeault cohomology. In this paper, we construct dGBV algebra for a general closed Kähler manifold. From this, we obtain a formal Frobenius manifold on the *A*-side for any closed Kähler manifold. Comparison with that of Barannikov-Kontsevich [1] suggests that these two kinds of formal Frobenius manifolds should be mirror image of each other. A standard argument in Kodaira-Spencer-Kuranishi deformation theory shows that one actually obtains a Frobenius (super)manifold structure on a neighborhood of the origin in the relevant cohomology, in the setting of both Barannikov-Kontsevich and ours. Now we have two formal Frobenius manifolds on the *A*-side: one from counting rational curves by Gromov-Witten invariants, the other from our construction. We conjecture that they can be identified. If this is true, it should have some applications in enumerative geometry.

Acknowledgements. *The authors would like to thank Gang Tian and S.-T. Yau for their interest. The work in this paper is carried out while the second author is visiting Texas A&M University. He likes to express his appreciation for the*

hospitality and financial support of the Mathematics Department and the Geometry-Analysis-Topology group. He also likes to thank Blaine Lawson for bringing his attention to the reference [12], which made him come across Barannikov-Kontsevich [1] and Manin [18] on the internet.

1. A CONSTRUCTION OF FROBENIUS (SUPER)MANIFOLDS

In this section, we review a construction of Frobenius supermanifolds. For details, the reader should consult the papers by Tian [23], Todorov [24], Barannikov-Kontsevich [1] and Manin [18]. Here, we follow the formulation by Manin [18].

The input of the construction is a differential Gerstenhaber-Batalin-Vilkovisky (dGBV) algebra (see e.g. [18], §5). Let $(\mathcal{A}, \wedge)$ be a supercommutative algebra with identity over a field $\mathbf{k}$, i.e., $(\mathcal{A}, \wedge)$ is an algebra with identity over $\mathbf{k}$, furthermore, $\mathcal{A}$ has $\mathbb{Z}_2$ -grading, such that for any homogeneous elements $a, b \in \mathcal{A}$ with degrees $|a|, |b|$ respectively, we have

$$a \wedge b = (-1)^{|a| \cdot |b|} b \wedge a.$$

Assume that there are two $\mathbf{k}$ -linear maps of odd degrees δ and Δ on $\mathcal{A}$, such that

- $\delta^2 = 0, \delta\Delta + \Delta\delta = 0, \Delta^2 = 0$.
- δ is a derivation: $\delta(a \wedge b) = (\delta a) \wedge b + (-1)^{|a|} a \wedge (\delta b)$.
- If $[a \bullet b] = (-1)^{|a|} (\Delta(a \wedge b) - \Delta a \wedge b - (-1)^{|a|} a \wedge \Delta b)$, then $[a \bullet \cdot] : \mathcal{A} \rightarrow \mathcal{A}$ is a derivation of degree $|a| + 1$, i.e.,

$$[a \bullet (b \wedge c)] = [a \bullet b] \wedge c + (-1)^{(|a|+1)|b|} b \wedge [a \bullet c].$$

The quintuple $(\mathcal{A}, \wedge, \delta, \Delta, [\bullet])$ with the above properties is called a dGBV algebra. As in Manin [18], §5, we have

$$\begin{aligned} [a \bullet b] &= -(-1)^{(|a|+1)(|b|+1)} [b \bullet a], \\ [a \bullet [b \bullet c]] &= [[a \bullet b] \bullet c] + (-1)^{(|a|+1)(|b|+1)} [b \bullet [a \bullet c]], \\ \Delta[a \bullet b] &= [\Delta a \bullet b] + (-1)^{|a|+1} [a \bullet \Delta b], \\ \delta[a \bullet b] &= [\delta a \bullet b] + (-1)^{|a|+1} [a \bullet \delta b]. \end{aligned}$$

We will be interested in the cohomology ring $H = H(\mathcal{A}, \delta)$, which we assume to be finite dimensional. An *integral* on $\mathcal{A}$ is an even linear functional $\int : \mathcal{A} \rightarrow \mathbf{k}$, such that

$$\begin{aligned} (1) \quad & \int (\delta a) \wedge b = (-1)^{|a|+1} \int a \wedge (\delta b), \\ (2) \quad & \int (\Delta a) \wedge b = (-1)^{|a|} \int a \wedge (\Delta b), \end{aligned}$$

for any homogeneous $a, b \in \mathcal{A}$. It follows from (1) that $\int$ induces a well-defined $\mathbf{k}$ -bilinear functional

$$\begin{aligned} (\cdot, \cdot) &: H(\mathcal{A}, \delta) \otimes H(\mathcal{A}, \delta) \rightarrow \mathbf{k}, \\ (\alpha, \beta) &= \int \alpha \wedge \beta, \end{aligned}$$

where $[\alpha]$ and $[\beta]$ are cohomology classes represented by δ -closed a and b respectively. If $(\cdot, \cdot)$ is nondegenerate, we say that the integral is *nice*. It is obvious that

$$(\alpha \wedge \beta, \gamma) = (\alpha, \beta \wedge \gamma).$$

By definition, $(H(\mathcal{A}, d), \wedge, (\cdot, \cdot))$ is then a *Frobenius algebra*, when $\mathcal{A}$ has a nice integral.

To find a deformation of the ring structure on H , consider $\delta_a : \mathcal{A} \rightarrow \mathcal{A}$ for even $a \in \mathcal{A}$. When a satisfies

$$(3) \quad \begin{aligned} \delta a + \frac{1}{2}[a \bullet a] &= 0, \\ \Delta a &= 0, \end{aligned}$$

then $(\mathcal{A}, \wedge, \delta_a, \Delta, [\bullet \bullet])$ is also a dGBV algebra. If $\int$ is an integral for $(\mathcal{A}, \wedge, \delta, \Delta, [\bullet \bullet])$, so is it for $(\mathcal{A}, \wedge, \delta_a, \Delta, [\bullet \bullet])$. If there is a natural way to identify $H(\mathcal{A}, \delta_a)$ with $H = H(\mathcal{A}, \delta)$ which preserves $(\cdot, \cdot)$, then we get another Frobenius algebra structure on H . The construction of Frobenius manifold structure is based on the existence of a solution $\Gamma = \sum \Gamma_n$ to (3), such that $\Gamma_0 = 0$, $\Gamma_1 = \sum x^j e_j$, $e_j \in \text{Ker } \delta \cap \text{Ker } \Delta$. For $n > 1$, $\Gamma_n \in \text{Im } \Delta$ is a homogeneous super polynomial of degree n in x^j 's, such that the total degree of Γ_n is even. Furthermore, x^0 only appears in Γ_1 . Such a solution is called a *normalized universal solution*. Its existence can be established inductively. This is how Tian [22] and Todorov [24] proved that the deformation of complex structures on a Calabi-Yau manifold is unobstructed. This was generalized by Barannikov-Kontsevich [1] to the case of extended moduli space of complex structures of a Calabi-Yau manifold. Manin [18] further generalized it to the case of dGBV algebras, which can be stated as follows:

Theorem 1.1. *Let $(\mathcal{A}, \wedge, \delta, \Delta, [\bullet \bullet])$ be a dGBV algebra which satisfies the following conditions:*

1. $H = H(\mathcal{A}, \delta)$ is finite dimensional.
2. There is a nice integral on $\mathcal{A}$.
3. The inclusions $i : (\text{Ker } \Delta, \delta) \hookrightarrow (\mathcal{A}, \delta)$ and $j : (\text{Ker } \delta, \Delta) \hookrightarrow (\mathcal{A}, \Delta)$ induce isomorphisms on cohomology.

Then there is a structure of formal Frobenius manifold on the formal spectrum of $\mathbf{k}[[H']]$, the algebra of formal power series generated by H' , where H' is the dual $\mathbf{k}$ -vector space of H .

2. FROBENIUS MANIFOLD STRUCTURES ON DOLBEAULT COHOMOLOGY

In this section, (X, g, J) will be a closed Kähler manifold with Kähler form ω . Consider the quadruple $(\Omega^{*,*}(X), \wedge, \delta = \bar{\partial}, \Delta = \partial^*)$. It is well-known that $\bar{\partial}^2 = 0$, $(\partial^*)^2 = 0$, and $\bar{\partial}\partial^* + \partial^*\bar{\partial} = 0$. Furthermore, $\bar{\partial}$ is a derivation. Set

$$[a \bullet b]_{\partial^*} = (-1)^{|a|}(\partial^*(a \wedge b) - \partial^*a \wedge b - (-1)^{|a|}a \wedge \partial b).$$

(To the authors' knowledge, a formula of this type in differential geometry was first discovered by Tian [22] (see also Todorov [24]) to prove the important result that deformations of Calabi-Yau manifolds are unobstructed.) Since both ω and J are parallel with respect to the Levi-Civita connection, it follows that near each $x \in X$, one can find a local frame $\{e^1, \dots, e^n\}$ of $T^{1,0}X$, such that $\omega = e^1 \wedge \bar{e}^1 + \dots + e^n \wedge \bar{e}^n$. Furthermore, $\nabla_{e_j} e_k = \nabla_{\bar{e}_j} e_k = \nabla_{e_j} \bar{e}_k = \nabla_{\bar{e}_j} \bar{e}_k = 0$ at x . Let $\{e^1, \dots, e^n\}$ be the dual basis. Then at x , for any $\alpha \in \Omega^{*,*}(X)$, we have (Griffiths-Harris [10], p. 113):

$$\begin{aligned} \partial \alpha &= e^j \wedge \nabla_{e_j} \alpha, & \bar{\partial} \alpha &= \bar{e}^j \wedge \nabla_{\bar{e}_j} \alpha, \\ \partial^* \alpha &= -e_j \lrcorner \nabla_{\bar{e}_j} \alpha, & \bar{\partial}^* \alpha &= -\bar{e}_j \lrcorner \nabla_{e_j} \alpha, \end{aligned}$$

Lemma 2.1. $[a \bullet (b \wedge c)]_{\partial^*} = [a \bullet b]_{\partial^*} \wedge c + (-1)^{(|a|+1)|b|} b \wedge [a \bullet c]_{\partial^*}.$

Proof. The left hand side is given by

$$\begin{aligned}
& (-1)^{|a|}(\partial^*(a \wedge b \wedge c) - (\partial^*a) \wedge b \wedge c - (-1)^{|a|}a \wedge \partial^*(b \wedge c)) \\
= & (-1)^{|a|}(-e_j \vdash \nabla_{\bar{e}_j}(a \wedge b \wedge c) + (e_j \vdash \nabla_{\bar{e}_j}a) \wedge b \wedge c \\
& + (-1)^{|a|}a \wedge e_j \vdash \nabla_{\bar{e}_j}(b \wedge c)) \\
= & (-1)^{|a|}(-e_j \vdash (\nabla_{\bar{e}_j}a \wedge (b \wedge c) + a \wedge \nabla_{\bar{e}_j}(b \wedge c)) + (e_j \vdash \nabla_{\bar{e}_j}a) \wedge b \wedge c \\
& + (-1)^{|a|}a \wedge e_j \vdash \nabla_{\bar{e}_j}(b \wedge c)) \\
= & (-1)^{|a|}(-(e_j \vdash \nabla_{\bar{e}_j}a) \wedge (b \wedge c) - (-1)^{|a|}\nabla_{\bar{e}_j}a \wedge e_j \vdash (b \wedge c) \\
& - (e_j \vdash a) \wedge \nabla_{\bar{e}_j}(b \wedge c) - (-1)^{|a|}a \wedge e_j \vdash \nabla_{\bar{e}_j}(b \wedge c) \\
& + (e_j \vdash \nabla_{\bar{e}_j}a) \wedge b \wedge c + (-1)^{|a|}a \wedge e_j \vdash \nabla_{\bar{e}_j}(b \wedge c)) \\
= & -\nabla_{\bar{e}_j}a \wedge e_j \vdash (b \wedge c) - (-1)^{|a|}(e_j \vdash a) \wedge \nabla_{\bar{e}_j}(b \wedge c).
\end{aligned}$$

The right hand side is computed in a similar fashion.

$$\begin{aligned}
& [a \bullet b] \wedge c + (-1)^{(|a|+1)|b|}b \wedge [a \bullet c] \\
= & (-1)^{|a|}(\partial^*(a \wedge b) - (\partial^*a) \wedge b - (-1)^{|a|}a \wedge \partial^*b) \wedge c \\
& + (-1)^{(|a|+1)|b|}(-1)^{|a|}b \wedge (\partial^*(a \wedge c) - (\partial^*a) \wedge c - (-1)^{|a|}a \wedge \partial^*c) \\
= & (-1)^{|a|}(-(e_j \vdash \nabla_{\bar{e}_j}(a \wedge b)) \wedge c + (e_j \vdash \nabla_{\bar{e}_j}a) \wedge b \wedge c \\
& + (-1)^{|a|}a \wedge (e_j \vdash \nabla_{\bar{e}_j}b) \wedge c) \\
& + (-1)^{(|a|+1)|b|}(-1)^{|a|}b \wedge (-e_j \vdash \nabla_{\bar{e}_j}(a \wedge c) + (e_j \vdash \nabla_{\bar{e}_j}a) \wedge c \\
& + (-1)^{|a|}a \wedge (e_j \vdash \nabla_{\bar{e}_j}c)) \\
= & -(-1)^{|a|}(e_j \vdash \nabla_{\bar{e}_j}(a \wedge b)) \wedge c - (-1)^{(|a|+1)|b|}(-1)^{|a|}b \wedge e_j \vdash \nabla_{\bar{e}_j}(a \wedge c) \\
& + a \wedge (e_j \vdash \nabla_{\bar{e}_j}b) \wedge c + (-1)^{|b|}a \wedge b \wedge (e_j \vdash \nabla_{\bar{e}_j}c) \\
& + 2(-1)^{|a|}(e_j \vdash \nabla_{\bar{e}_j}a) \wedge b \wedge c \\
= & -\nabla_{\bar{e}_j}a \wedge (e_j \vdash b) \wedge c - (-1)^{|b|}\nabla_{\bar{e}_j}a \wedge b \wedge (e_j \vdash c) \\
& - (-1)^{|a|}(e_j \vdash a) \wedge \nabla_{\bar{e}_j}b \wedge c - (-1)^{|a|}(e_j \vdash a) \wedge b \wedge \nabla_{\bar{e}_j}c \\
= & -\nabla_{\bar{e}_j}a \wedge e_j \vdash (b \wedge c) - (-1)^{|a|}(e_j \vdash a) \wedge \nabla_{\bar{e}_j}(b \wedge c).
\end{aligned}$$

□

Thus $(\Omega^{*,*}(X), \wedge, \delta = \bar{\partial}, \Delta = \partial^*)$ is a dGBV algebra.

Remark 2.1. One can obtain an indirect proof as follows. Cao-Zhou [6] showed that for a Kähler manifold, $\sqrt{-1}(\bar{\partial}^* - \partial^*)$ is the operator Δ defined in Koszul [13], who also defined the *covariant Schouten-Nijenhuis bracket*

$$\{a, b\} = (\Delta a) \wedge b + (-1)^{|a|}a \wedge (\Delta b) - \Delta(a \wedge b)$$

which satisfies the property

$$\{a, b \wedge c\} = \{a, b\} \wedge c + (-1)^{(|a|+1)|b|}b \wedge \{a, c\}.$$

As pointed out in Cao-Zhou [6],

$$[a \bullet b] = (-1)^{|a|}\{a, b\}$$

is the bracket generated by Δ . Using the type decomposition and $[a \bullet (b \wedge c)] = [a \bullet b] \wedge c + (-1)^{(|a|+1)|b|} b \wedge [a \bullet c]$, one then gets

$$[a \bullet (b \wedge c)]_{\partial^*} = [a \bullet b]_{\partial^*} \wedge c + (-1)^{(|a|+1)|b|} b \wedge [a \bullet c]_{\partial^*}.$$

Let $\int_X : \Omega^{*,*}(X) \rightarrow \mathbb{C}$ be the ordinary integration of differential forms. Analogous to Claim 4.1 in Barannikov-Kontsevich [1], we have

Lemma 2.2. *The following identity holds on closed Kähler manifolds:*

$$\begin{aligned} \int_X \bar{\partial} a \wedge b &= (-1)^{|a|+1} \int_X a \wedge \bar{\partial} b, \\ \int_X \partial^* a \wedge b &= (-1)^{|a|} \int_X a \wedge \partial^* b, \end{aligned}$$

Proof. By linearity, we can assume that a has type (p, q) . To prove the first identity, we only need to consider the case of b having type $(n-p, n-q-1)$. Then we have $\partial(a \wedge b) = 0$, since it has type $(n+1, n-1)$. Hence $d(a \wedge b) = \bar{\partial}(a \wedge b)$. Similarly, we have $da \wedge b = \bar{\partial} a \wedge b$. By Stokes theorem, we have

$$\int_X \bar{\partial} a \wedge b + (-1)^{|a|} a \wedge \bar{\partial} b = \int_X \bar{\partial}(a \wedge b) = \int_X d(a \wedge b) = 0.$$

To prove the second equality, recall that $\partial^* = - * \partial *$, and $*^2 \alpha = (-1)^{|\alpha|} \alpha$ (see Griffiths-Harris [10], p. 82). Then since $|b| = |a| - 1$, we have

$$\begin{aligned} \int_X \partial^* a \wedge b &= (-1)^{|b|} \int_X \partial^* a \wedge (*b) \\ &= (-1)^{|b|} (\partial^* a, *b) = (-1)^{|b|} (a, \partial^* b) \\ &= (-1)^{|b|} \int_X a \wedge * \partial^* b = -(-1)^{|b|} \int_X a \wedge \partial^* b \\ &= (-1)^{|a|} \int_X a \wedge \partial^* b. \end{aligned}$$

□

Proposition 2.1. *The two natural inclusions $i : (\text{Ker } \partial^*, \bar{\partial}) \rightarrow (\Omega^{*,*}(X), \bar{\partial})$ and $j : (\text{Ker } \bar{\partial}, \partial^*) \rightarrow (\Omega^{*,*}(X), \partial^*)$ induce isomorphisms on cohomology.*

Proof. Consider the Laplace operators $\Delta_{\partial} = \partial \partial^* + \partial^* \partial$ and $\Delta_{\bar{\partial}} = \bar{\partial} \bar{\partial}^* + \bar{\partial}^* \bar{\partial}$. On Kähler manifold (see e.g. Griffiths-Harris [10], p. 115), we have $\Delta_{\partial} = \Delta_{\bar{\partial}} = \frac{1}{2} \Delta_d$. By Hodge theory, every element in $H(\Omega^{*,*}, \bar{\partial})$ is represented by an element α , such that $\Delta_{\bar{\partial}} \alpha = 0$. It then follows that $\Delta_{\partial} \alpha = 0$, and hence $\partial^* \alpha = 0$, i.e., $\alpha \in \text{Ker } \partial^*$. This proves that $H(i)$ is surjective.

To show that $H(i)$ is injective, let $\alpha \in \text{Ker } \partial^*$, such that $\alpha = \bar{\partial} \beta$ for some β . We need to show that $\alpha \in \bar{\partial} \text{Ker } \partial^*$. By Hodge decomposition, $\bar{\partial} \beta = \Delta_{\bar{\partial}} \gamma$ for some γ . Then we have

$$\Delta_{\bar{\partial}} \bar{\partial} \gamma = \bar{\partial} \Delta_{\bar{\partial}} \gamma = \bar{\partial}^2 \gamma = 0,$$

which implies that $\bar{\partial} \gamma = 0$. Furthermore,

$$\Delta_{\partial} \partial^* \gamma = \partial^* \Delta_{\partial} \gamma = \partial^* \Delta_{\bar{\partial}} \gamma = \partial^* \alpha = 0$$

implies that $\partial^* \gamma = 0$. Now $\alpha = \Delta_{\bar{\partial}} \gamma = \bar{\partial}(\bar{\partial}^* \gamma) + \bar{\partial}^*(\bar{\partial} \gamma) = \bar{\partial}(\bar{\partial}^* \gamma)$. But

$$\partial^*(\bar{\partial}^* \gamma) = -\bar{\partial}^*(\partial^* \gamma) = 0$$

show that $\bar{\partial}^* \gamma \in \text{Ker } \partial^*$, therefore, $\alpha \in \bar{\partial} \text{Ker } \partial^*$, thus $H(i)$ is injective. The proof for $H(j)$ is similar. □

Now we use Theorem 1.1 to get

Theorem 2.1. *For any closed Kähler manifold X , if K is the algebra of formal power series generated by the dual space of the Dolbeault cohomology $H_{\bar{\partial}}^*(X)$, then there is a structure of formal Frobenius manifold on the formal spectrum of K obtained from the dGBV algebra $(\Omega^{*,*}(X), \wedge, \delta = \bar{\partial}, \Delta = \partial^*)$.*

Remark 2.2. The construction involves the solution by power series method to the equation

$$\bar{\partial}\Gamma + \frac{1}{2}[\Gamma, \Gamma] = 0,$$

where $\Gamma = \sum_n \Gamma_n$, $\Gamma_0 = 0$, $\Gamma_1 = \sum x^j \eta_j$, Γ_n is a (super)polynomial in x^j 's with coefficients in $\text{Im } \bar{\partial}\partial^*$. Here η_j is a basis of $\text{Ker } \bar{\partial} \cap \text{Ker } \partial^* \cong H = H(\Gamma(X, \Lambda^* TX \otimes \Lambda^* \bar{T}^* X), \bar{\partial})$, x^j is the dual basis of H' . Notice that Γ is a (super)polynomial for x^j 's (Fermionic coordinates) with odd degrees, and is a power series only for x^j 's with even degrees (Bosonic coordinates). A modification of the standard argument in Kodaira-Spencer-Kuranishi deformation theory (see e.g. Morrow-Kodaira [20], §4.2) shows that these series converges for all Fermionic x^j 's, and for small values of Bosonic x^j 's (this method was also used by Tian [22] and Todorov [24]). Therefore, one gets a Frobenius supermanifold with the Bosonic part an open neighborhood of the zero vector in H^{even} .

Remark 2.3. The same construction carries through if we take $\delta = \partial$, $\Delta = \bar{\partial}^*$.

3. COMPARISON WITH BARANNIKOV-KONTSEVICH'S FROBENIUS MANIFOLD

Assume that X is a closed Calabi-Yau n -manifold, fix a nowhere vanishing holomorphic n -form $\Omega \in \Gamma(X, \Lambda^n T^* X)$. Barannikov-Kontsevich [1] considered the dGBV algebra with

$$\mathcal{A} = \oplus \mathcal{A}_k, \quad \mathcal{A}_k = \oplus_{q+p=k} \Gamma(X, \Lambda^p TX \otimes \Lambda^q \bar{T}^* X).$$

Here $\mathcal{A}$ is $\mathbb{Z}$ -graded. It also has an induced $\mathbb{Z}_2$ -grading. This grading is different from that in [1] (shifted by 1), we adopt this grading to be compatible with the notations of Manin [18]. The multiplication $\wedge$ on $\mathcal{A}$ is given by the ordinary wedge products on $\Lambda^* \bar{T}^* X$ and $\Lambda^* TX$. The derivation is $\delta = \bar{\partial}$. Notice that for any two integers $0 \leq p, q \leq n$, Ω defines an isomorphism

$$\Lambda^p TX \otimes \Lambda^q \bar{T}^* X \rightarrow \Lambda^{n-p} T^* X \otimes \Lambda^q \bar{T}^* X, \quad \gamma \mapsto \gamma \lrcorner \Omega,$$

defined by contraction of the p -vector with Ω to get a form of type $(n-p, 0)$. Then Δ is defined by

$$(\Delta\gamma) \lrcorner \Omega = \partial(\gamma \lrcorner \Omega).$$

Tian-Todorov lemma shows that the bracket, defined by

$$[\alpha \bullet \beta] = (-1)^{|\alpha|} (\Delta(\alpha \wedge \beta) - \Delta\alpha \wedge \beta - (-1)^{|\alpha|} \alpha \wedge \Delta\beta),$$

is given by the wedge product on type $(0, *)$ -forms and the Schouten-Nijenhuis bracket on type $(*, 0)$ polyvector fields. It is straightforward to check that $(\mathcal{A}, \wedge, \delta, \Delta)$ as above is a dGBV algebra. Furthermore, the linear functional

$$\int : \mathcal{A} \rightarrow \mathbb{C}, \quad \int \gamma = \int_X (\gamma \lrcorner \Omega) \wedge \Omega$$

is a nice integral on X .

Since the complex $(\mathcal{A}, \delta)$ contains the deformation complex

$$\Omega^{0,0}(TX) \xrightarrow{\bar{\partial}} \Omega^{0,1}(TX) \xrightarrow{\bar{\partial}} \Omega^{0,2}(TX),$$

It is called the extended deformation complex. Barannikov-Kontsevich [1] (Lemma 2.1) showed that the deformation functor associated with the graded differential Lie algebra $(\mathcal{A}[-1], \delta, [\cdot \bullet \cdot])$ is represented by the formal spectrum of formal power series generated by H' , the dual of $H = H(\mathcal{A}, \delta)$. This is the extended moduli space of complex structures in Witten [28]. Furthermore, they constructed a structure of a formal Frobenius manifold on it.

Remark 3.1. As in Remark 2.2, One gets Frobenius supermanifold structure.

To facilitate the comparison with our construction, we now rewrite the operator Δ in Barannikov-Kontsevich [1], which appeared in earlier work of Tian [23] and Todorov [24] with different notation, in terms of more familiar operator. To start with, notice that the Hermitian metric induces an isomorphism $TX \cong \overline{T}^*X$. Therefore, we get, for any two integers $0 \leq p, q \leq n$, an isomorphism

$$f_{p,q} : \Lambda^p TX \otimes \Lambda^q \overline{T}^*X \rightarrow \Lambda^p \overline{T}^*X \otimes \Lambda^q TX.$$

On $\Lambda^p \overline{T}^*X \otimes \Lambda^q TX$, we consider the operator

$$\bar{\partial}^* : \Gamma(X, \Lambda^p \overline{T}^*X \otimes \Lambda^q TX) \rightarrow \Gamma(X, \Lambda^{p-1} \overline{T}^*X \otimes \Lambda^q TX).$$

Lemma 3.1. *We have the following commutative diagram:*

$$\begin{array}{ccc} \Gamma(X, \Lambda^p TX \otimes \Lambda^q \overline{T}^*X) & \xrightarrow{\Delta} & \Gamma(X, \Lambda^{p-1} TX \otimes \Lambda^q \overline{T}^*X) \\ f_{p,q} \downarrow & & \downarrow f_{p-1,q} \\ \Gamma(X, \Lambda^p \overline{T}^*X \otimes \Lambda^q TX) & \xrightarrow{-\bar{\partial}^*} & \Gamma(X, \Lambda^{p-1} \overline{T}^*X \otimes \Lambda^q TX). \end{array}$$

Proof. It is well-known that both ω and Ω are parallel with respect to the Levi-Civita connection. It follows that near each $x \in X$, one can find a local frame $\{e^1, \dots, e^n\}$ of $T^{1,0}X$, such that $\Omega = e^1 \wedge \dots \wedge e^n$, $\omega = e^1 \wedge \bar{e}^1 + \dots + e^n \wedge \bar{e}^n$. Furthermore, $\nabla_{e_j} e_k = \nabla_{\bar{e}_j} e_k = \nabla_{e_j} \bar{e}_k = \nabla_{\bar{e}_j} \bar{e}_k = 0$ at x . Let $\{e^1, \dots, e^n\}$ be the dual basis. Near x , every $\gamma \in \Gamma(X, \Lambda^{p-1} TX \otimes \Lambda^q \overline{T}^*X)$ can be written as

$$\gamma = \sum \gamma_{JK} e_J \otimes \bar{e}^K,$$

where J runs over multiple indices $(j_1, \dots, j_p)$, $1 \leq j_1 < \dots < j_p \leq n$, K run over multiple indices $(k_1, \dots, k_q)$, $1 \leq k_1 < \dots < k_q \leq n$, $e_J = e_{j_1} \wedge \dots \wedge e_{j_p}$, $\bar{e}^K = \bar{e}^{k_1} \wedge \dots \wedge \bar{e}^{k_q}$. Then we have

$$\begin{aligned} f_{p-1,q}^{-1} \bar{\partial}^* f_{p,q} \gamma &= f_{p-1,q}^{-1} \left(\sum \bar{\partial}^* (\gamma_{JK} \bar{e}^J \wedge e_K) \right) \\ &= -f_{p-1,q}^{-1} \left(\sum (\nabla_{e_j} \gamma_{JK}) (\bar{e}_j \lrcorner \bar{e}^J) \wedge e_K \right) \\ &= -\sum (\nabla_{e_j} \gamma_{JK}) (e^j \lrcorner e_J) \wedge \bar{e}^K. \end{aligned}$$

On the other hand,

$$\begin{aligned} (\Delta \gamma) \lrcorner \Omega &= \partial(\gamma \lrcorner \Omega) \\ &= \sum \partial(\gamma_{JK} (e_J \lrcorner \Omega) \wedge \bar{e}^K) \\ &= \sum (\nabla_{e_j} \gamma_{JK}) (e^j \wedge (e_J \lrcorner \Omega)) \wedge \bar{e}^K. \end{aligned}$$

It is then obvious that $f_{p-1,q}^{-1} \bar{\partial}^* f_{p,q} \gamma = -\Delta \gamma$. $\square$

Denote $\widehat{\partial}^* = f_{p-1,q}^{-1} \bar{\partial}^* f_{p,q}$. We have the following interesting comparison of the two dGBV algebras:

$$(BK) \quad \mathcal{A} = \oplus_{p,q} \Gamma(X, \Lambda^p T X \otimes \Lambda^q \overline{T}^* X), \quad \delta = \bar{\partial}, \quad \Delta = -\widehat{\partial}^*,$$

$$(CZ) \quad \mathcal{A} = \oplus_{p,q} \Gamma(X, \Lambda^p T^* X \otimes \Lambda^q \overline{T}^* X), \quad \delta = \bar{\partial}, \quad \Delta = \partial^*,$$

(The definitions of the multiplications are similar!) It is reasonable to conjecture that the two corresponding Frobenius manifolds are mirror images of each other.

REFERENCES

- [1] S. Barannikov, M. Kontsevich, *Frobenius Manifolds and Formality of Lie Algebras of Polyvector Fields*, preprint, alg-geom/9710032.
- [2] J. Bertin, C. Peters, *Variations de structures de Hodge, variétés de Calabi-Yau et symétrie miroir*, in **Introduction à la théorie de Hodge**, 169–256, Panor. Synthèses, 3, Soc. Math. France, Paris, 1996.
- [3] J.-L. Brylinski, *A differential complex for Poisson manifolds*, **J. Differential Geom.** **28** (1988), no. 1, 93–114.
- [4] P. Candelas, X. C. de la Ossa, p. S. Green, L. Parkes, *A pair of Calabi-Yau manifolds as an exactly soluble superconformal theory*, **Nuclear Phys. B** **359** (1991), no. 1, 21–74.
- [5] H.-D. Cao, J. Zhou, *On Quantum de Rham Cohomology Theory*, preprint, April, 1998.
- [6] H.-D. Cao, J. Zhou, *Identification of Two Frobenius Manifolds In Mirror Symmetry*, preprint, 1998.
- [7] B. Dubrovin, *Integrable systems in topological field theory*, **Nuclear Phys. B** **379** (1992), no. 3, 627–689.
- [8] B. Dubrovin, *Geometry of 2D topological field theories*, in **Integrable systems and quantum groups** (Montecatini Terme, 1993), 120–348, Lecture Notes in Math., 1620, Springer, Berlin, 1996.
- [9] W.M. Goldman, J.J. Millson, *The deformation theory of representations of fundamental groups of compact Kähler manifolds*, **Inst. Hautes études Sci. Publ. Math. No. 67** (1988), 43–96.
- [10] P. Griffiths, J. Harris, *Principles of algebraic geometry*. Pure and Applied Mathematics. Wiley-Interscience [John Wiley & Sons], New York, 1978.
- [11] M. Kontsevich, *Homological algebra of mirror symmetry*, in **Proceedings of the International Congress of Mathematicians**, Vol. 1, 2 (Zürich, 1994), 120–139, Birkhäuser, Basel, 1995.
- [12] M. Kontsevich, *Deformation quantization of Poisson manifolds, I*, preprint, q-alg/9709040.
- [13] J.-L. Koszul, *Crochet de Schouten-Nijenhuis et cohomologie*, **The mathematical heritage of Élie Cartan (Lyon, 1984)**, *Astisque* 1985, Numéro Hors Serie, 257–271.
- [14] J. Li, G. Tian, *Comparison of the algebraic and the symplectic Gromov-Witten invariants*, preprint, alg-geom/9712035.
- [15] B. Lian, K. Liu, S.T. Yau, *Mirror principle I*, preprint, alg-geom/9712011.
- [16] G. Liu, *Associativity of quantum multiplication*, preprint, 1994.
- [17] D. McDuff, D. Salamon, *J-holomorphic curves and quantum cohomology*. University Lecture Series, 6. American Mathematical Society, Providence, RI, 1994.
- [18] Y. Manin, *Three constructions of Frobenius manifolds: a comparative study*, preprint, math.QA/9801006.
- [19] D.R. Morrison, *Mirror symmetry and rational curves on quintic threefolds: a guide for mathematicians*, **J. Amer. Math. Soc.** **6** (1993), no. 1, 223–247.
- [20] J. Morrow, K. Kodaira **Complex manifolds**, Holt, Rinehart and Winston, Inc., New York-Montreal, Que.-London, 1971.
- [21] Y. Ruan, G. Tian, *A mathematical theory of quantum cohomology*, **J. Differential Geom.** **42** (1995), no. 2, 259–367.
- [22] G. Tian, *Smoothness of the universal deformation space of compact Calabi-Yau manifolds and its Petersson-Weil metric*, in **Mathematical aspects of string theory** (San Diego, Calif., 1986), 629–646, Adv. Ser. Math. Phys., 1, World Sci. Publishing, Singapore, 1987.

- [23] G. Tian, *Quantum cohomology and its associativity*, in **Current developments in mathematics**, 1995 (Cambridge, MA), 361–401, Internat. Press, Cambridge, MA, 1994.
- [24] A.N. Todorov, *The Weil-Petersson geometry of the moduli space of $SU(n \geq 3)$ (Calabi-Yau) manifolds. I.*, **Comm. Math. Phys.** **126** (1989), no. 2, 325–346.
- [25] C. Vafa, *Topological mirrors and quantum rings*, in **Essays on mirror manifolds**, S.T. Yau ed., 96–119, Internat. Press, Hong Kong, 1992.
- [26] I. Vaisman, **Lectures on the geometry of Poisson manifolds**. Progress in Mathematics, 118. Birkhuser Verlag, Basel, 1994.
- [27] E. Witten, *Two-dimensional gravity and intersection theory on moduli space*, in **Surveys in differential geometry** (Cambridge, MA, 1990), 243–310, Lehigh Univ., Bethlehem, PA, 1991.
- [28] E. Witten, *Mirror manifolds and topological field theory*, in **Essays on mirror manifolds**, S. T. Yau ed., International Press Co., Hong Kong, 1992.
- [29] S.T. Yau ed., **Essays on mirror manifolds**, International Press, Hong Kong, 1992.

DEPARTMENT OF MATHEMATICS, TEXAS A & M UNIVERSITY, COLLEGE STATION, TX 77843
E-mail address: cao@math.tamu.edu, zhou@math.tamu.edu